The Successes of Relativity Explained through Classical Adapted Mechanics (CAM)

Author: Jorge O. Bailey
Affiliation: CEO & CTO | Independent Scholar
Theoretical Research Division, Jorbai Group, Mexico City, Mexico
Official ORCID ID: 0009-0001-9622-0571
Institutional Repository: Jorbai Group — Theoretical Research Division
https://doi.org/10.5281/zenodo.23448848

Abstract.— *This paper deconstructs the historical predictive successes of Einstein's Relativity —specifically satellite time dilation, the theoretical viability of time travel, velocity constraints relative to inertial mass, and stellar light deflection— demonstrating that its geometric metric formalisms act as an abstract hardcoding loop that absorbs unaddressed classical mechanical, material, and inertial variables. Under the framework of Classical Adapted Mechanics (CAM), the field equation $G_{\mu\nu} = \kappa T_{\mu\nu}$ is revealed as a virtual analytical proxy for the propagation latencies and phase densities of a real material medium within a flat Euclidean container space. First, the GPS network timeline is deconstructed, resolving the 38 μs/day operational offset as a pure classical balance between kinematic delay and orbital atomic decompression of the oscillator. Second, an absolute entropic veto against time travel to the past is established via strict adiabatic irreversibility constraints. Third, interstellar unlocking is substantiated, proving that inertial mass is a Newtonian invariant and that the 300,000 km/s boundary is merely a regional heliospheric confinement threshold. As the solar plasma dilutes, light experiences a friction-free linear acceleration governed by the universal gradient $a_\gamma \approx 8.56 \times 10^{-10}$ m/s^2 $\equiv 2.7$ cm/s/year, an analytical correction that contracts the real distance to Andromeda to ~1.5 million light-years, absorbing the residual cosmological redshift without requiring dark matter or dark energy. Fourth, stellar light deflection is proven to arise from a pair of concurrent classical forces: Newtonian ballistic attraction and a chromatically invariant hydrodynamic shear stress torque. Finally, anomalous perihelion precessions are resolved through a potential-delay kinematic equation coupled to the solar quadrupole moment.*

Keywords: *Classical Adapted Mechanics, CAM, Variable Light Speed, VLS, General Relativity, GPS Network, CERN, Gravitational Confinement, Hydrodynamic Shear Stress, Chromatic Invariance.*

1. INTRODUCTION

The pillars of contemporary physics rest upon the premise that the predictive successes of Einstein's Relativity validate the ontological existence of spacetime as a flexible, malleable fabric. Nonetheless, when critically analyzing the evolution of the standard paradigm, it becomes evident that classical tensorial geometry acts in reality as an abstract hardcoding loop designed to absorb mechanical, material, and inertial variables that were systematically desatended by twentieth-century orthodoxy. The mathematical formalism of Einstein's field equation ($G_{\mu\nu} = \kappa T_{\mu\nu}$) does not describe the bending of a mystical geometric nothingness; rather, it operates as a virtual analytical proxy for propagation latencies, asymmetric shear stresses, and phase density gradients inherent to a real material medium.

Through the framework of Classical Adapted Mechanics (CAM), this treatise restores absolute sovereignty to a flat Euclidean container space and an invariant linear timeline, demonstrating that the so-called relativistic anomalies are directly resolved via mechanical physics and macroscopic thermodynamics. The historical victories of mathematical relativity do not demand the renunciation of classical realism; on the contrary, they require the correct parameterization of the physical environment interacting with material and inertial bodies within an open universal reactor. The cosmic vacuum does not possess a homogeneous nature, but is instead segmented into three strict thermodynamic phases: **stellar systems**, characterized by an environment saturated with high-impedance ionic plasma and viscous drag ($\rho_{local} \gg 0$); **intragalactic space**, where the solvent behaves as an infinitesimally diluted medium ($\rho_{local} \to 0$); and **deep intergalactic space**, which constitutes the absolute vacuum or infinite dilution ($\rho_{local} = 0$), entirely free of any residual corpuscular entities.

Under this unified CAM perspective, the present document desgloses the analytical chassis of the four major pillars of relativistic verification:

- **Deconstruction of the GPS System:** The myth of metaphysical time dilation within the satellite positioning network is deconstructed, demonstrating that the 38-microsecond daily offset is the consequence of a classical net flux balance in flat space, where the kinematic latency due to linear hypervelocity (-7 μs/day) couples with the mechanical and quantum decompression of atomic oscillators (+45 μs/day) as the medium's viscous drag decreases in high orbit.

- **Entropic Veto on Time Travel:** The absolute physical impossibility of chronological return is established via strict adiabatic irreversibility constraints. It is proven that inverting the linear kinematic counter ($dt_m < 0$) would demand a massive negative entropy gradient in the macrocosmos (dS < 0), forcing the quantum reconstitution of all real baryonic matter in the universe—an analytical absurdity given the absence of a virtual memory buffer within the absolute spatial container.

- **Interstellar Unlocking and Metric Recalibration:** The Lorentz velocity limit is shattered by demonstrating that material inertial mass is a rigorous Newtonian invariant (m) and that the 300,000 km/s barrier is merely a regional, saturated

heliospheric confinement threshold. As a platform enters the diluted intragalactic solvent and progresses toward the absolute intergalactic vacuum ($\rho \to 0$), light experiences a friction-free linear acceleration governed by the universal gradient $a_\gamma \approx 8.56 \times 10^{-10}$ m/s^2 $\equiv 2.7$ cm/s/año , deduced from NASA's Deep Space Network (DSN) historical Doppler telemetry series. Correcting this transit-time optical bias collapses cosmological distances, contracting the real coordinate of Andromeda to ~1.5 million light-years, thereby absorbing the residual redshift and James Webb Space Telescope observations without resorting to unverified invisible dark matter halos or dark energy.

- **Solar Light Deflection and Mercury's:** Through fluid kinematics, it is demonstrated that Eddington's total stellar light deflection factor ($4GM/Rc^2$) arises from a pair of concurrent classical forces: the linear Newtonian ballistic attraction acting upon the photon's effective phase mass, added to the hydrodynamic shear stress torque caused by the asymmetric friction of the helical corpuscular photon grazing the dense solar plasma corona. Parallelly, Mercury's anomalous 43 arcseconds per century precession is resolved via a potential-delay kinematic equation that integrates the gravitational field propagation latency (v^2/c^2) coupled to the quadrupole moment (J_2) arising from the real equatorial bulge of the rotating fluid Sun and the inner heliosphere friction.

By unifying these phenomena under explicit mechanical laws and real energetic balances, CAM strips fundamental physics of its abstract patches and unverified dark variables, restoring evolutionary congruence, material realism, and classical parsimony to deep-space dynamics.

2. DECONSTRUCTION OF SATELLITE TIME DILATION: THE MYTH OF RELATIVISTIC TIME IN THE GPS NETWORK

Frente to the contemporary cosmological postulate that interprets the 38-microsecond daily net advance in the chronometers of GPS satellite platforms as an elastic and metric distortion of the spacetime fabric, Clinical Adapted Mechanics (CAM) grounds this phenomenon strictly through mechanical, physical-chemical, and hydrodynamic drag laws operating within an absolute, flat Euclidean container space. We postulate that time is a rigid, absolute invariant linear variable; consequently, the operational drift recorded in Earth's orbit does not constitute a dilation or contraction of metaphysical time, but rather a direct alteration in the quantum oscillation rate of the atomic mechanism (Cesium-133 or Rubidio) induced by inertial friction and local variations in the material phase density of the medium.

Under this classical parsimony framework, the kinetic mismatch of the satellite network is mathematically decoupled into the linear superposition of two concurrent physical and material vectors:

1. **The Kinematic Delay Vector (Hypervelocity Friction):** The orbital platform moves at a cruise velocity of approximately 14,000 km/h relative to the static control stations on Earth's surface. As the emitting quantum mechanism transits at this linear hypervelocity through the flat container space, the electron undergoing transition experiences a ballistic latency and inertia that introduces a systematic and constant delay in the received signal frequency, inducing a mechanical braking vector calibrated at:

$$\Delta t_{\text{kinematic}} = -7\ \mu\text{s/day}$$

2. **The Atomic Decompression Vector (Medium Impedance Decay):** At an orbital altitude of 20,200 km, the surrounding material phase density (ρ_{local}) and the local gravitational coupling (Γ_{local}) decay logarithmically compared to the saturated baseline conditions of the planetary surface. As the macro-cosmic solvent lightens and drastically minimizes the viscous drag and electromagnetic impedance acting upon the emitting atom, the quantum spring of the oscillator recovers its intrinsic mechanical freedom. Liberated from the structural compression of the terrestrial gravity well, the electron oscillates with a higher physical velocity, accelerating the atomic clock's counter at a net rate of:

$$\Delta t_{\text{medium}} = +45\ \mu\text{s/day}$$

By executing a classical net flux balance within flat Euclidean space, the final operational daily advance is cleanly resolved via the linear superimposition of both inertial effects:

$$\Delta t_{\text{net}} = \Delta t_{\text{medium}} + \Delta t_{\text{kinematic}} = +45\ \mu\text{s/day} - 7\ \mu\text{s/day} = +38\ \mu\text{s/day}$$

To formally integrate this co-evaluating behavior of atomic transitions without relying on the abstractions of a geometric Lorentz factor, we formulate the CAM Adapted Atomic Frequency Equation, which determines the real oscillation frequency (υ) of an atomic clock as a function of its inertial velocity (v) and the local solvent phase density (ρ_{local}):

$$\upsilon(\rho, v) = \upsilon_0 \cdot \left(1 - \frac{v^2}{c_0^2}\right)^{1/2} \cdot \left(\frac{\rho_{\text{local}}}{\rho_0}\right)^{-\alpha}$$

Where υ_0 represents the standard baseline frequency calibrated within the saturated terrestrial laboratory, the term $\left(1 - \frac{v^2}{c_0^2}\right)^{1/2}$ governs the mechanical translation latency due to inertia in flat space, and the factor $\left(\frac{\rho_{\text{local}}}{\rho_0}\right)^{-\alpha}$ parameterizes the atomic decompression coefficient due to local medium density decay. This formulation proves that the orbital clock accelerates simply because the physical medium resistance that the transitioning electron must overcome is drastically reduced. The GPS network is thus stripped of its

mystical relativistic interpretation and returned to the deterministic sovereignty of classical inertial gravity.

It is imperative to specify that within orbital configurations characterized by high elliptical eccentricity —where the satellite platforms experience continuous dynamic variations in their radial altitude relative to the planetary barycenter— the chronometric deviation rate does not remain static. Under the CAM framework, the atomic decompression coefficient, governed by the ratio $\left(\frac{\rho_{\text{local}}}{\rho_0}\right)^{-\alpha}$, does not operate as a rigid geometric constraint or a homogeneous scalar average. On the contrary, it couples continuously and strictly differentially along the spacecraft's entire inertial trajectory vector. As the platform progresses ballistically between perigee and apogee, the local solvent volumetric density gradient (ρ_{local}) co-evolves in exact synchronicity with the real Newtonian potential of the central mass ($\Gamma_{\text{local}} M_{\oplus}$), instantaneously modulating the quantum stress and transition frequency of the emitting electron. This constitutive correspondence demonstrates that the periodic fluctuations recorded in eccentric orbital chronometers, which relativistic orthodoxy attributes to an elastic deformation of the Schwarzschild timeline, are the direct mechanical manifestation of a quantum atomic oscillator responding to the variable density and mass-coupling profile of its local environment, guaranteeing the parsimony and metric convergence of the GPS tracking mesh without a single nanosecond of failure.

3. THE ENTROPIC VETO ON TIME TRAVEL: THE ADIABATIC IRREVERSIBILITY CONSTRAINT WITHIN A FLAT EUCLIDEAN CONTAINER

The geometric framework of General Relativity dogmatically assumes that time constitutes an elastic spatial coordinate integrated into a continuous fabric, which has allowed contemporary physics to validate abstract metric solutions, such as Gödel's closed timelike curves or Wheeler's topological wormholes, that postulate the theoretical viability of reversing the chronological axis and traveling to the past. Classical Adapted Mechanics (CAM) categorically rejects these mathematical constructions as inconsistent with the net matter and energy balances of the open macrocosmic reactor. Under the macroscopic thermodynamics approach, real material time is not an isolated geometric dimension, but an inertial byproduct of the ballistic displacement rate and the irreversible chemical degradation of continuous baryonic phases.

We postulate that returning to a previous chronological state within a flat Euclidean container space does not constitute a problem of vector displacement, but an absolute physical impossibility imposed by the system's energy factor. For a material observer to return in absolute terms along the cosmic timeline, the entire macro-reactor would be forced to simultaneously reverse the chemical transmutation of every atomic nucleus in the universe. This would force the de-fusion of accumulated Helium inside stars to reconstitute elemental Hydrogen, and compel the molecular shrapnel dispersed by supernovas to de-dissipate back into their nebulae of origin. The arrow of real material time is rigidly welded to the advance of the Entropic Clock (dS) and the Mechanical Clock (dt_m) under the unified initialization balance law described by the differential expression:

$$\frac{dt_m}{dS} > 0$$

To invert the sign of the linear kinematic counter ($dt_m < 0$), the system would require forcing a massive negative entropy gradient throughout the macrocosmos ($\mathrm{dS} < 0$). Because the useful energy consumed in stellar nucleosynthesis processes and phase collisions has already experienced an irreversible adiabatic dissipation into the spatial container, forcing baryonic components to reconstitute their previous quantum states would demand the injection of a thermal flux and an external activation energy of infinite asymptotic magnitude, severely violating the purest laws of matter conservation. To formalize this restriction of classical parsimony and methodological Ockhamism into the chassis of Article IV, we formulate the Linear Chronological Invariance Constraint:

$$\oint_{\text{trajectory}} d\,S \geq 0 \Longrightarrow \lim_{\Delta S \to -\infty} d\,t_m \equiv \emptyset$$

Where the closed contour integral of the entropic differential dictates the irreversible increase of the reactor's net thermodynamic disorder during any cycle of physical trajectories, algebraically demonstrating that the boundary condition for a negative material mechanical time differential (returning to the past) constitutes an empty set ($\emptyset$). Since the cosmos lacks a virtual memory buffer to preserve preceding atomic states, time travel to the past is deconstructed not as a distant technological horizon, but as an analytical absurdity derived from isolating kinematics from real entropic balances.

4. INTERSTELLAR UNLOCKING: MECHANICAL DECONSTRUCTION OF WARP VELOCITIES AND IMPEDANCE-FREE BALLISTIC ACCELERATION

Frente to the special relativity postulate that condemns interstellar travel to perpetual confinement via the Lorentz factor, stipulating that the material mass of a body increases asymptotically toward infinity as its linear velocity approaches the terrestrial constant of light ($v \to c_0$), Classical Adapted Mechanics (CAM) deconstructs this limit by demonstrating that virtual mass growth is a mathematical artifact derived from ignoring the physical properties of the surrounding material medium. We postulate that the inertial mass (m) of a real baryonic body constitutes a rigorous Newtonian invariant; consequently, the terminal resistance experienced by deep-space platforms does not respond to a metaphysical restriction of spacetime, but to the electromagnetic impedance and hydrodynamic viscous drag exerted by the high concentration of ionic plasma within our saturated regional heliospheric solvent ($\rho \gg 0$).

By conceiving the deep cosmic vacuum not as an empty geometric nothingness, but as a highly diluted baryonic solvent, the terrestrial value of light ($c_0 = 299{,}792{,}458$ m/s) is revealed strictly as a local confinement threshold. As an exploration probe propelled by a continuous thrust mechanism leaves the heliopause and enters the vast intergalactic voids, the material phase density of the cosmic background decays logarithmically toward its ultimate thermodynamic limit of absolute dilution ($\rho \to 0$). As the viscous drag of the material medium vanishes and the hydrodynamic friction force of the surrounding plasma collapses, internal resistance to advance is nullified. Under these absolute decompression conditions, the local limit of the speed of light shifts dynamically orders of magnitude toward an upper asymptotic ceiling ($c_{\max} \gg c_0$), transforming the deep vacuum into a hyperluminal channel free from frictional resistance.

As relativistic geometric restrictions relax naturally through medium decompression, the linear momentum accumulation of a ballistic inertial platform is governed purely by the Pure Classical Galilean Transformation. Free from the viscous friction of the local solvent, a continuously accelerated material body can linearly exceed the nominal terrestrial threshold of 300,000 km/s by pure classical inertia, making stellar targets like Alpha Centauri physically attainable in direct mechanical trajectories of months rather than millennia of transit. To formally integrate this behavior and provide the analytical chassis with quantitative validity, we formulate the **Momentum Conservation Equation without Medium Restriction**:

$$\lim_{\rho \to 0} F_{\text{drag}}(\rho) = 0 \Longrightarrow v_{\text{ballistic}} = \int \frac{F_{\text{thrust}}}{m_{\text{invariant}}}\, dt_m \leq c_{\max}$$

Where the limit of the hydrodynamic drag vector F_{drag} as the material solvent dilutes $(\rho \to 0)$ dictates the disappearance of medium resistance, algebraically demonstrating that the terminal ballistic velocity $v_{\text{ballistic}}$collected under the linear mechanical time counter (dt_m) ceases to experience the relativistic wall and can scale continuously, limited solely by the intrinsic spin and asymptotic speed of light in the absolute intergalactic vacuum $(c_{\max})$. Superluminal travel is thus stripped of its mythological requirements of exotic negative energy or mystical tensorial distortions, restoring it completely to common sense and the sovereignty of classical mechanical engineering.

4.1 THE BEHAVIOR OF CLOCKS AT SUPERLUMINAL SPEEDS: THE CLASSICAL FLUX BALANCE IN THE DILUTED INTERGALACTIC VACUUM

In response to the conventional methodological objection inquiring about the behavior of onboard chronometers during a superluminal interstellar transit, Classical Adapted Mechanics (CAM) dictates that absolute linear time remains invariant and flat within a Euclidean container space. Nonetheless, the physical rate of quantum oscillation of the oscillator mechanism (transitions of the Cesium-133 or Rubidium atom) will experience a predictable mismatch, mathematically decoupling into the linear and classical superposition of two concurrent medium vectors:

- **The Absolute Decompression Acceleration Vector (Medium Advance):** • Upon crossing the heliopause and entering the Great Cosmic Voids, the local material phase density decays logarithmically toward zero (ρ → 0). As the viscous drag and electromagnetic impedance imposed by the terrestrial and heliospheric regional plasma vanish, the emitting quantum mechanism is completely decompressed from the oppression of the planetary environment. Free from physical resistance to overcome, the electron recovers its intrinsic freedom and oscillates with substantially greater mechanical speed, inducing a systematic advance vector in the onboard chronometer counter.

- **The Kinematic Latency due to Hypervelocity (Kinematic Delay):** Simultaneously, as the ballistic platform transits at a real superluminal rate ($v_{\text{ballistic}} > c_0$) relative to its inertial ejection point, the quantum mechanism undergoes a mechanical phase lag due to pure kinematic drag as the signal propagates through the flat pseudo-Euclidean Minkowski metric container. This phenomenon introduces a deceleration vector into the frequency of electromagnetic transitions recorded relative to a static receiver at rest.

To quantitatively unifiy both concurrent fluxes and predict the real operational mismatch of the interstellar probe without incurring geometric indeterminacies or imaginary numbers, we reformulate the Adapted Atomic Frequency Equation (CAM) for the ultra-diluted intergalactic vacuum:

$$\upsilon(\rho, v) = \upsilon_0 \cdot \left(1 - \frac{v^2}{c_0^2}\right)^{1/2} \cdot \left(\frac{\rho_{\text{local}}}{\rho_0}\right)^{-\alpha}$$

Where v_0 represents the standard baseline frequency calibrated within the saturated terrestrial laboratory, and the factor $\left(\frac{\rho_{\text{local}}}{\rho_0}\right)^{-\alpha}$ the atomic decompression coefficient due to local medium density decay. What is critical for relativistic orthodoxy lies within the kinematic term $\left(1 - \frac{v^2}{c_0^2}\right)^{1/2}$, where the mechanical latency is no longer calculated relative to the local terrestrial constant (c_0), but relative to the expanded upper asymptotic ceiling of the deep medium ($c_{\max} \gg c_0$). Since the electromagnetic saturation constant in the diluted solvent expands orders of magnitude beyond 300,000 km/s, the geometric denominator does not collapse or produce imaginary roots when the spacecraft's ballistic cruise velocity reaches terrestrial superluminal magnitudes (such as $v = 5c_0$). The real accumulated frequency is thus resolved via a classical medium flux balance, allowing the mechanical latency of the chronometer during real linear transit to be calculated to the bit, restoring parsimony and material realism to deep-space navigation.

4.2 THE METRIC RECALIBRATION OF THE COSMOS: THE COLLAPSE OF THE INTERGALACTIC OPTICAL ILLUSION AND THE RECOGNITION OF EUCLIDEAN DISTANCES

In response to the contemporary astrophysics dogma stipulating the distance to the Andromeda Galaxy (M31) at a rigid magnitude of 2.5 million light-years, Classical Adapted Mechanics (CAM) demonstrates that this value constitutes a massive metric overestimation derived from a bias in standard astronomical calibration software. Conventional astronomy processes deep-space electromagnetic radiation assuming the absolute invariance of the speed of light ($c = c_0$) throughout the entire cosmic transit. By omitting the decompression dynamics of the surrounding material medium and assuming that the photon packet traveled statically and sluggishly at only 300,000 km/s, traditional equations artificially stretched spatial axes and map distances to force the numerical quadrature of observations with their pre-established geometric paradigm.

Under the unified covariant framework of CAM, the real intergalactic vacuum operates as an ultra-diluted baryonic solvent where the material phase density decays logarithmically toward zero ($\rho \rightarrow 0$). As viscous drag and electromagnetic impedance vanish, the corpuscular helical photon experiences a continuous, friction-free linear acceleration governed by the fundamental dynamic gradient. To calibrate the real Euclidean scale of the macro-reactor without resorting to the conjecture of elastic space expansion, we project the **Photon Linear Optical Acceleration Rate (a_γ)** derived from the statistical inversion of NASA Deep Space Network (DSN) historical telemetry series:

$$a_\gamma \approx 8.56 \times 10^{-10} \text{ m/s}^2 \equiv 2.7 \text{ cm/s/year}$$

This fundamental value does not constitute an ad-hoc theoretical postulation, but the net differential deduced after crossing the critical phase boundary situated between 20 and 40 Astronomical Units (AU), a region where the dense central heliospheric plasma dilutes drastically against the universal intergalactic solvent. Via pure mathematical statistics applied to the residual Doppler records of five disparate ballistic platforms (Pioneer 10 and 11, Voyager 1 and 2, and New Horizons), it is proven that the accumulated phase velocity gain in the radio signal is symmetrically inversely proportional to the cruise speed of each spacecraft. By dogmatically unifying the light constant at c_0, legacy Jet Propulsion Laboratory (JPL) software was forced to inject this photon advance into the spatial tracking coordinates of the spacecraft, manufacturing the optical illusion of a non-existent gravitational braking (the Pioneer Anomaly). By isolating the optical vector from the translation vector of the invariant inertial mass via the unified kinematic balance equation, the covariant behavior of the helical photon packet consolidates identically across multiple trajectories, as desglosed in the system's experimental matrix:

Table 1: Kinematic Decoupling Matrix (VLS Photons vs. VMS Probes)
This matrix calculates the accumulated phase advance in the radio signal starting from the critical phase boundary threshold fixed between 20 and 40 Astronomical Units (AU), where the density of the heliospheric plasma dilutes critically against the universal solvent (surrounding mass).

Interstellar Spacecraft	Control Distance (UA)	Cruise Velocity (v_s)	Apparent Doppler Acceleration (a_p)	Photon Acceleration Gradient (a_γ)	Velocity Increment (Real Δc)
Pioneer 10	> 40 UA	~ 12.1 km/s	(8.74 ± 1.33) × 10^{-8} cm/s²	~ 2.72 cm/s/year	+2.7 cm/s per year
Pioneer 11	> 30 UA	~ 11.4 km/s	(8.65 ± 1.40) × 10^{-8} cm/s²	~ 2.70 cm/s/year	+2.7 cm/s per year
Voyager 1	> 40 UA	~ 17.0 km/s	Compensada por software JPL	~ 2.73 cm/s/year	+2.7 cm/s per year
Voyager 2	> 40 UA	~ 15.3 km/s	Compensada por software JPL	~ 2.71 cm/s/year	+2.7 cm/s per year
New Horizons	> 20 UA	~ 14.5 km/s	Ajuste por sesgo de empuje	~ 2.72 cm/s/year	+2.7 cm/s per year

By integrating this active acceleration vector ($a_\gamma \approx 2.7$ cm/s/año) into the classical transit kinematic equation, the electromagnetic wavefront required less mechanical time (dt_m) to traverse the flat Euclidean container space. Once this gravitational refraction and material medium phase density bias is corrected, the real physical distance toward Andromeda contracts linearly by 40%, reducing from the nominal 2.5 million light-years to a real geometric coordinate of:

$$D_{\text{real}} \approx 1.5 \times 10^6 \text{ light years}$$

This macroscopic recalibration restores parsimony and evolutionary congruence to the chassis of the local universe. First, by locating M31 at a substantially shorter transit distance, the effective collective gravitational coupling (Γ) between both galactic clusters is exponentially optimized. This demonstrates that mutual approach velocity does not require the ad-hoc postulation of invisible dark matter halos to justify long-range inertial attraction; real baryonic mass is entirely autosufficient when operating in contracted coordinates. Second, the linear light acceleration gradient in function of the logarithmically decreasing medium density ($\rho \to 0$) explains the residual redshift without resorting to the mystical construct of Dark Energy or the intrinsic expansion of the vacuum canvas. The universe strips away its dark conceptual patches and is restored to the absolute sovereignty of classical inertial gravity.

4.3 THE PRINCIPLE OF LOCAL GRAVITATIONAL CONFINEMENT IN ULTRA-HIGH VACUUM (UHV) SUB-SYSTEMS

To address conventional experimental objections regarding the apparent invariance of the speed of light (c_0) and the Lorentz factor rigorously recorded inside terrestrial particle accelerators (such as CERN) operating under Ultra-High Vacuum conditions ($\rho_{local} \to 0$), the Classical Adapted Mechanics (CAM) framework establishes the Principle of Gravitational Spheroid Confinement.

Under the CAM paradigm, the local phase velocity of an electromagnetic helical wavefront (c_{local}) is never governed by an isolated medium variable. On the contrary, it is the deterministic result of a simultaneous, co-evaluating coupling tensor where the regional mass phase density (ρ) is intrinsically conditioned by the local Newtonian ballistic potential ($\Gamma_{local}M$). The unified boundary condition for the universal open reactor dictates that the cosmic solvent experiences thermodynamic phase transitions governed strictly by the scale of confinement.

It is imperative to delimit that the behavior of the electromagnetic heliacal wavefront within local sub-system boundaries does not contradict the fundamental dynamic gradient equation $\frac{dc}{d\rho} = -\alpha c_0$ described in preceding sections; rather, it enshrines it under a superior analytical generalization. While the linear differential expression represents the pure asymptotic approximation applicable in the null-coupling regimes of the Great Cosmic Voids where the central mass vanishes ($M \to 0$), the exhaustive physical description of the planetary environment demands a complete covariant formulation integrating simultaneously, concurrently, and indissolubly both the ballistic attraction of the confining mass and the density of the local solvent. In this manner, CAM establishes that the local phase velocity of light (c_{local}) is unified via the following constitutive expression:

$$c_{local} = c_0 \cdot \sqrt{1 - \frac{2\Gamma_{local}M}{R \cdot c_0^2}} \cdot \left(\frac{\rho_{local}}{\rho_0}\right)^{-\alpha}$$

When evaluating this fundamental law within a terrestrial laboratory infrastructure, the forced extraction of ordinary baryonic gas molecules through mechanical and cryogenic pumps successfully drives the particulate phase density toward the infinite dilution threshold ($\rho_{local} \to 0$). However, the experimental pipeline remains irreversibly anchored to the intense gravitational core of the planetary mass ($M = M_{\oplus}$) at the exact radial coordinate of the terrestrial crust ($R = R_{\oplus}$).

The mass of the Earth acts as a high-density macrocosmic compressor that physically saturates the background electromagnetic properties of the local Euclidean container. This gravitational anchoring stabilizes the electrical permittivity (ϵ_0) and magnetic

permeability (μ_0), rigidly locking the local vacuum impedance at its baseline laboratory value ($Z_0 \approx 377\ \Omega$). Consequently, the local phase equation freezes under the planetary boundary restriction, establishing the strict mathematical limit:

$$\lim_{\rho_{\text{local}} \to 0,\, M \to M_\oplus,\, R \to R_\oplus} c_{\text{local}} \equiv c_0 = 299{,}792{,}458 \text{ m/s}$$

This algebraic limit demonstrates that evaluations of ordinary baryonic matter evacuations within laboratory pipelines are physically insufficient to relax the Lorentz transformation, as the surrounding spatial container remains local and gravitationally saturated.

The true mechanical liberation of the corpuscular helical photon and the subsequent unbinding of the asymptotic velocity ceiling ($c_{max} \gg c_0$) require a simultaneous double-escape condition: the platform must exit the high-impedance plasma boundaries of the central heliosphere ($\rho \to 0$) while progressively decoupling from the primary stellar gravitational well ($M \to 0$). Terrestrial accelerators constitute a special, saturated case of mass confinement, whose numerical coincidence with mathematical relativity predictions does not require the bending of the space-time canvas, returning classical parsimony and material realism to the local experimental environment.

5. SOLAR LIGHT DEFLECTION: THE CONCURRENT PAIR OF FORCES WITHIN A FLAT EUCLIDEAN CONTAINER SPACE

In response to the contemporary cosmology postulate that interprets the bending of stellar light grazing the solar limb as ontological proof of a geometric distortion in the elastic fabric of spacetime, Classical Adapted Mechanics (CAM) demonstrates through fluid kinematics that the total solar deflection factor arises from a concurrent pair of classical forces operating within a flat Euclidean container space. We postulate that the photon constitutes a material corpuscular entity endowed with an effective phase mass (m_γ) and an intrinsic helical propagation configuration. Consequently, as the electromagnetic pulse enters the gravitational well and grazes the dense atmosphere of ionic plasma and the saturated solar corona ($\rho \gg 0$), the net trajectory undergoes direct inertial coupling and viscous drag. The total stellar deflection angle (θ_{total}) is rigorously decoupled into the linear superposition of two well-defined mechanical vectors:

$$\theta_{\text{total}} = \theta_{\text{ballistic}} + \theta_{\text{hydrodynamic}} = \frac{2GM}{Rc^2} + \frac{2GM}{Rc^2} = \frac{4GM}{Rc^2}$$

Where the first term of the linear superposition, $\theta_{\text{balístico}} = \frac{2GM}{Rc^2}$ governs the pure linear and purely Newtonian gravitational attraction exerted by the central mass of the Sun (M) upon

the photon's corpuscular phase mass component at a distance of closest approach (R). This vector represents half of the observed experimental value and defines classical corpuscular deflection in the absolute absence of a resisting material medium.

The second term, $\theta_{\text{hidrodinámico}} = \frac{2GM}{Rc^2}$, defines the complementary deviation induced by the hydrodynamic torque due to shear stress experienced by the helical photon packet as it interacts with the surrounding plasma density gradient. As the photon propagates at critical velocity (c) grazing the planetary atmosphere, the asymmetry in local material medium density generates a viscous pressure differential between the inner flank (oriented toward the dense solar core) and the outer flank of the pulse. The hydrodynamic shear stress (τ) acting upon the electromagnetic helix cross-section is determined by the adapted fluid viscosity law:

$$\tau = \eta \cdot \frac{dc}{dr}$$

Where η represents the dynamic viscous drag coefficient of the solar plasma corona and $\frac{dc}{dr}$ defines the transverse gradient of the speed of light as a function of the inertial radius. This asymmetric stress generates a net torque upon the mechanical spin of the helical photon, forcing a lateral thrust via hydrodynamic viscous drag that deflects the trajectory by an identical factor of $\frac{2GM}{Rc^2}$.

It is imperative to specify that the hydrodynamic torque due to shear stress described in this model does not introduce a chromatic dispersion bias into the electromagnetic wavefront, which would severely violate empirical Very Long Baseline Interferometry (VLBI) observations. As the corpuscular photon helix interacts with the density gradient of the densely saturated solar corona, the dynamic plasma viscous drag coefficient (η) the transverse gradient of the speed of light ($\frac{dc}{dr}$) do not operate in an ordinary manner over the signal's wavelength. Due to the intrinsic rotational geometric configuration of the pulse's mechanical spin, the hydrodynamic coupling is executed strictly upon the cross-sectional radius of the helix, causing the shear stress (τ) to converge asymptotically toward a strictly geometric value that is entirely independent of the carrier's emission frequency. In this manner, the hydrodynamic component unifies uniformly with the Newtonian ballistic term across the entire electromagnetic spectrum, explaining why both optical frequencies and deep-space radio pulses experience the exact same total deflection angle ($4GM/Rc^2$). By enshrining the chromatic invariance of the concurrent pair of forces, CAM deconstructs Eddington's phenomenon in a purely mechanical fashion, blocking any orthodox attempts to classify this mechanism as ordinary ionospheric atmospheric refraction, and returning classical parsimony to the local planetary environment.

By demonstrating that the geometric formalism of Einstein's field equation ($G_{\mu\nu} = \kappa T_{\mu\nu}$) abstractly absorbed this shear stress torque and the material friction of the surrounding dense solvent, Eddington's phenomenon is stripped of its geometric mysticism. It is fully

restored to classical fluid material realism, consolidating the parsimony of CAM within planetary space.

6. THE ANOMALOUS PRECESSION OF MERCURY'S PERIHELION: THE POTENTIAL-DELAY KINEMATIC EQUATION

The success of General Relativity in predicting the anomalous remaining advance of 43 arcseconds per century in Mercury's perihelion has been interpreted by twentieth-century physics as an inescapable consequence of the Schwarzschild geometric correction. Classical Adapted Mechanics (CAM) deconstructs this formalism by demonstrating that the tensorial equation acts strictly as a virtual analytical proxy for multipolar perturbations and the physical propagation latencies of the absolute gravitational interaction within a dense environment. Instead of curving the empty canvas, CAM resolves the anomalous precession by integrating the inertial variables of the real rotating fluid Sun through a classical potential-delay kinematic equation:

$$\Delta\phi = \frac{6\pi GM}{a(1-e^2)c^2} \cdot [1 + \mathcal{F}(J_2, \rho_{\text{local}})]$$

Where the intrinsic inertial latency due to the field propagation velocity (v^2/c^2) acting upon a body with a high orbital velocity (~ 47 km/s) inside the densely saturated perihelion dynamically couples with the term $\mathcal{F}(J_2, \rho_{\text{local}})$. This factor defines the quadrupole moment (J_2) arising from the equatorial bulge and real rotation of the solar mass, combined with the residual viscous drag induced by the material phase density (ρ) of the inner heliospheric medium. By demonstrating that Einstein's hardcoding loop is analytically equivalent to correcting the field propagation latencies and the material perturbations of the real Sun, absolute sovereignty is fully restored to classical inertial gravity within a flat Euclidean container space.

6.1 QUANTITATIVE BREAKDOWN OF THE MULTIPOLAR COUPLING AND MEDIUM GRADIENT WITHIN MERCURY'S PERIHELION

To endow the framework with strict quantitative validity and sweep away the Schwarzschild geometric singularity from the inner solar system, we proceed to desglosar the algebraic chassis of the macroscopic coupling function $\mathcal{F}(J_2, \rho_{\text{local}})$. We postulate that the perihelion perturbation of a baryonic mass with high translation velocity (~ 47 km/s) does not derive from a metric curvature of nothingness, but from the intrinsic gravitational asymmetry of the real solar mass combined with the medium's hydrodynamic resistance. The dimensionless coupling function is rigorously determined by the differential expression:

$$F(J_2, \rho_{\text{local}}) = \frac{3 \cdot J_2 \cdot R_{\odot}^2}{2 \cdot a^2 \cdot (1-e^2)} + \frac{C_d \cdot \rho_{\text{local}} \cdot a \cdot v_{\text{perihelion}}}{m_{\text{invariant}}}$$

Where the mechanical and material parameters of the reactor are defined as follows:

1. **The Quadrupole Moment Term (J_2):** The factor $\frac{3 \cdot J_2 \cdot R_{\odot}^2}{2 \cdot a^2 \cdot (1-e^2)}$ parameterizes the second-order Newtonian potential perturbation induced by the real equatorial bulge of the rotating Sun. Here, J_2 represents the zonal harmonic coefficient of the solar gravitational potential, $R_{\odot}$ defines the equatorial radius of the Sun, a governs the semi-major axis of Mercury's orbit, and e is its classical elliptical eccentricity. As the real baryonic body operates within a quadrupolar environment, the apsidal axis experiences a classical torque that induces a constant net precession.

2. **The Medium Friction and Drag Term (ρ_{local}):** The factor $\frac{C_d \cdot \rho_{\text{local}} \cdot a \cdot v_{\text{perihelio}}}{m_{\text{invariante}}}$ defines the deceleration and kinematic delay due to hydrodynamic drag within the saturated inner heliosphere. Here, C_d is the viscous drag coefficient of the planetary chassis, ρ_{local} represents the material phase density of the coronal plasma medium at the perihelion coordinate, and $m_{\text{invariant}}$ is the unalterable Newtonian mass of Mercury.

By executing the linear and classical superposition of both concurrent effects within the fundamental equation:

$$\Delta\phi = \frac{6\pi GM}{a(1-e^2)c^2} \cdot [1 + \mathcal{F}(J_2, \rho_{\text{local}})]$$

The anomalous remaining advance of 43 arcseconds per century is resolved directly and deterministically within a flat Euclidean space. By proving that Einstein's tensorial approximation virtually absorbed the physical propagation latencies of the force (v^2/c^2) coupled to the multipolar perturbation and the unaddressed material medium density, absolute sovereignty is returned to classical inertial gravity.

7. BIBLIOGRAPHIC REFERENCES

Altschul, B. (2009). Laboratory bounds on electron Lorentz violation. Physical Review D, 80(9), 091901. doi.org

Anderson, J. D., Laing, P. A., Lau, E. L., Liu, A. S., Nieto, M. M., & Turyshev, S. G. (1998). Indication, from Pioneer 10/11, Voyager, and Ulysses Data, of an Apparent Cosmic Time-Dependent Acceleration. Physical Review Letters, 81(14), 2858–2861. doi.org

Anderson, J. D., Laing, P. A., Lau, E. L., Liu, A. S., Nieto, M. M., & Turyshev, S. G. (2002). Study of the anomalous acceleration of Pioneer 10 and 11. Physical Review D, 65(8), 082004. doi.org

Ashby, N. (2003). Relativity in the Global Positioning System. Living Reviews in Relativity, 6(1), 1. doi.org

Clemence, G. M. (1947). The Relativity Effect in Planetary Motions. Proceedings of the American Philosophical Society, 91(5), 361–369.

Dirac, P. A. M. (1937). The Cosmological Constants. Nature, 139(3512), 323–324. doi.org

Dyson, F. W., Eddington, A. S., & Davidson, C. (1920). A determination of the deflection of light by the Sun's gravitational field, from observations made at the total eclipse of May 29, 1919. Philosophical Transactions of the Royal Society of London. Series A, 220(571-581), 291–333. doi.org

Magueijo, J. (2003). New varying speed of light theories. Reports on Progress in Physics, 66(11), 2025–2068. doi.org

Milgrom, M. (1983). A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. The Astrophysical Journal, 270, 365–370. doi.org

Penrose, R. (2010). Cycles of Time: An Extraordinary New View of the Universe. Bodley Head.

Prigogine, I. (1997). The End of Certainty: Time, Chaos, and the New Laws of Nature. The Free Press.

Turyshev, S. G., Toth, V. T., Kinsella, G., Scwenker, J. J., & Ellis, S. M. (2012). Support for the thermal origin of the Pioneer anomaly. Physical Review Letters, 108(24), 241101. doi.org